# Author Index


Advenier, C., see Moreau, J. 576(1992)103
Aggarwal, S. K., Kinter, M. and Herold, D. A.
Determination of cobalt in urine by gas chromatography–mass spectrometry employing nickel as an internal standard 576(1992)297
Alvarez-Bujidos, M. L., see Negro, A. 576(1992)135
Arai, I., see Shibasaki, H. 576(1992)47
Ardemagni, R., see Lombardi, F. 576(1992)129
Aucoin, D. P., see Tyczkowska, K. L. 576(1992)305
Bagchi, D., see Shara, M. A. 576(1992)221
Banno, K.
Analytical studies on the chiral separation and simultaneous determination of pantothenic acid and hopantenic acid enantiomers in rat plasma by gas chromatography–mass fragmentography. Reply to Küsters *et al.* 576(1992)181
Barham, C., see Liu, H. 576(1992)163
Baudendistel, L. J., see Scott, M. D. 576(1992)149
Beijnen, J. H., see Van Tellingen, O. 576(1992)158
Bellato, P., see Mariot, R. 576(1992)143
Boison, J. O., Korsrud, G. O., MacNeil, J. D., Keng, L. and Papich, M.
Determination of penicillin G in bovine plasma by high-performance liquid chromatography after pre-column derivatization 576(1992)315
Brondz, I. and Olsen, I.
Intra-injector methylation of free fatty acids from aerobically and anaerobically cultured *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 576(1992)328
Bucheli, F., see Wyss, R. 576(1992)111
Caspary, W. F., see Stein, J. 576(1992)53
Chen, C.-F., see Tsai, T.-H. 576(1992)170
Chen, I.-W., Lin, J. H. and Dorley, J.
Quantitation of a new cholecystokinin and gastrin receptor antagonist (L-365,260) in dog and rat plasma by high-performance liquid chromatography 576(1992)363
Chen, S., see Mariot, R. 576(1992)143
Colzani, V., see Lombardi, F. 576(1992)129
Cordonnier, P., see Moreau, J. 576(1992)103
Cubría, J. C., see Negro, A. 576(1992)135
Dahms, T. E., see Scott, M. D. 576(1992)149
Davies, J. H., see Poon, G. K. 576(1992)235
Delgado, M., see Liu, H. 576(1992)163
De Smet, M., see Van Belle, S. J.-P. 576(1992)351
Dickson, P. H., see Shara, M. A. 576(1992)221
Dorley, J., see Chen, I.-W. 576(1992)363
Eder, C., see Küsters, E. 576(1992)179
Eggers, C., see Liu, H. 576(1992)163
Favarato, M., Mizzen, C. A. and McLachlan, D. R.
Resolution of serum aluminum-binding proteins by size-exclusion chromatography: identification of a new carrier of aluminum in human serum 576(1992)271
Forman, L. J., see Liu, H. 576(1992)163
Francotte, E. and Junker-Buchheit, A.
Preparative chromatographic separation of enantiomers (Review) 6248(1992)1
Fuchs, M. J. and Keim, V.
Separation of rat pancreatic secretory proteins by cation-exchange fast protein liquid chromatography 576(1992)287
Fukuoka, H., see Takahara, E. 576(1992)174
Furuta, T., Katayama, M., Shibasaki, H. and Kasuya, Y.
Simultaneous determination of stable isotopically labelled L-histidine and urocanic acid in human plasma by stable isotope dilution mass spectrometry 576(1992)213
Furuta, T., see Shibasaki, H. 576(1992)47
Geerts, F., see Van Belle, S. J.-P. 576(1992)351
Gupta, R. N.
Drug level monitoring: antidepressants (Review) 576(1992)183
Hawkes, W. C. and Kutnink, M. A.
High-performance liquid chromatographic determination of selenocysteine with the fluorescent reagent, N-(iodoacetylaminoethyl)-5-naphthylamine-1-sulfonic acid 576(1992)263
Heeremans, C. E. M., see Poon, G. K. 576(1992)235
Herold, D. A., see Aggarwal, S. K. 576(1992)297
Horowitz, J. D., see Sallustio, B. C. 576(1992)321
Jarman, M., see Poon, G. K. 576(1992)235
Junker-Buchheit, A., see Francotte, E. 6248(1992)1
Kasuya, Y., see Furuta, T. 576(1992)213
Kasuya, Y., see Shibasaki, H. 576(1992)47
Katayama, M., see Furuta, T. 576(1992)213
Kato, H., see Takahara, E. 576(1992)174
Kawakami, Y., Ohga, T., Shimamoto, C., Satoh, N. and Ohmori, S.
Determination of free N-acetylamino acids in biological samples and N-terminal acetylamino acids of proteins 576(1992)63
Kawamoto, M., see Morita, I. 576(1992)334
Keim, V., see Fuchs, M. J. 576(1992)287
Keng, L., see Boison, J. O. 576(1992)315
Kermici, M., see Krien, P. M. 576(1992)255
Kinter, M., see Aggarwal, S. K. 576(1992)297
Korsrud, G. O., see Boison, J. O. 576(1992)315
Krien, P. M., Margou, V. and Kermici, M.
Electrochemical determination of femtomole amounts of free reduced and oxidized glutathione. Application to human hair follicles 576(1992)255
Kulemeier, J., see Stein, J. 576(1992)53
Küsters, E., Spöndlin, C. and Eder, C.
Analytical studies on the chiral separation and simultaneous determination of pantothenic acid and hopantenic acid enantiomers in rat plasma by gas chromatography–mass fragmentography: a reply 576(1992)179
Kutnink, M. A., see Hawkes, W. C. 576(1992)263

Lagendijk, J., Ubbink, J. B. and Vermaak, W. J. H.
Quantification of erythrocyte S-adenosyl-L-methionine levels and its application in enzyme studies 576(1992)95
Larsson, L., see Magnusson, P. 576(1992)79
Laufen, H., see Riedel, K.-D. 576(1992)358
Lembcke, B., see Stein, J. 576(1992)53
Leung, P. Y. and Tsao, C. S.
Preparation of an optimum mobile phase for the simultaneous determination of neurochemicals in mouse brain tissues by high-performance liquid chromatography with electrochemical detection 576(1992)245
Liau, L. S., see Ong, C.-N. 576(1992)346
Lin, J. H., see Chen, I.-W. 576(1992)363
Liu, H., Forman, L. J., Montoya, J., Eggers, C., Barham, C. and Delgado, M.
Determination of valproic acid by high-performance liquid chromatography with photodiode-array and fluorescence detection 576(1992)163
Löfman, O., see Magnusson, P. 576(1992)79
Lombardi, F., Ardemagni, R., Colzani, V. and Visconti, M.
High-performance liquid chromatographic determination of rufloxacin and its main active metabolite in biological fluids 576(1992)129
MacNeil, J. D., see Boison, J. O. 576(1992)315
Magnusson, P., Löfman, O. and Larsson, L.
Determination of alkaline phosphatase isoenzymes in serum by high-performance liquid chromatography with post-column reaction detection 576(1992)79
Margou, V., see Krien, P. M. 576(1992)255
Mariot, R., Zangirolami, L., Bellato, P., Chen, S., Pieraccini, G. and Moneti, G.
Determination of the coumarin derivative cloricromene acid in rabbit plasma and platelets 576(1992)143
Massart, D. L., see Van Belle, S. J.-P. 576(1992)351
McCague, R., see Poon, G. K. 576(1992)235
McLachlan, D. R., see Favarato, M. 576(1992)271
Méndez, R., see Negro, A. 576(1992)135
Michalski, W. P.
Resolution of three forms of superoxide dismutase by immobilised metal affinity chromatography 576(1992)340
Milligan, P. A.
Determination of piroxicam and its major metabolites in the plasma, urine and bile of humans by high-performance liquid chromatography 576(1992)121
Mizzen, C. A., see Favarato, M. 576(1992)271
Moneti, G., see Mariot, R. 576(1992)143
Monsaert, C., see Van Belle, S. J.-P. 576(1992)351
Montoya, J., see Liu, H. 576(1992)163
Moreau, J., Palette, C., Cordonnier, P., Naline, E., Advenier, C. and Pays, M.
Separation and identification of the 4-hydroxyantipyrine sulphoconjugate 576(1992)103
Morita, I., Kawamoto, M. and Yoshida, H.
Difference in the concentration of tryptophan metabolites between maternal and umbilical foetal blood 576(1992)334
Morris, R. G., see Sallustio, B. C. 576(1992)321
Nagasawa, T., see Sakai, T. 576(1992)155
Nagata, O., see Takahara, E. 576(1992)174
Naline, E., see Moreau, J. 576(1992)103
Negro, A., Alvarez-Bujidos, M. L., Ortiz, A. I., Cubría, J. C., Méndez, R. and Ordóñez, D.
Reversed-phase ion-pair high-performance liquid chromatographic determination of triclabendazole metabolites in serum and urine 576(1992)135
Niessen, W. M. A., see Poon, G. K. 576(1992)235
Nooyen, W. J., see Van Tellingen, O. 576(1992)158
Ohga, T., see Kawakami, Y. 576(1992)63
Ohmori, S., see Kawakami, Y. 576(1992)63
Olsen, I., see Brondz, I. 576(1992)328
Ong, C.-N., Liau, L. S. and Ong, H. Y.
Separation of globins using free zone capillary electrophoresis 576(1992)346
Ong, H. Y., see Ong, C.-N. 576(1992)346
Ordóñez, D., see Negro, A. 576(1992)135
Ortiz, A. I., see Negro, A. 576(1992)135
Palette, C., see Moreau, J. 576(1992)103
Papich, M., see Boison, J. O. 576(1992)315
Patin, E., see Rivat, C. 576(1992)71
Pays, M., see Moreau, J. 576(1992)103
Pieraccini, G., see Mariot, R. 576(1992)143
Poon, G. K., Jarman, M., McCague, R., Davies, J. H., Heeremans, C. E. M., Van der Hoeven, R. A. M., Niessen, W. M. A. and Van der Greef, J.
Identification of 4-hydroxyandrost-4-ene-3,17-dione metabolites in prostatic cancer patients by liquid chromatography–mass spectrometry 576(1992)235
Regnault, V., Rivat, C., Vallar, L. and Stoltz, J.-F.
Dye-affinity purification of transthyretin from an unexploited by-product of human plasma chromatographic fractionation 576(1992)87
Riedel, K.-D., Wildfeuer, A., Laufen, H. and Zimmermann, T.
Equivalence of a high-performance liquid chromatographic assay and a bioassay of azithromycin in human serum samples 576(1992)358
Rivat, C., see Regnault, V. 576(1992)87
Rivat, C., Sertillanges, P., Patin, E. and Stoltz, J. F.
Single-step method for purification of human transferrin from a by-product of chromatographic fractionation of plasma 576(1992)71
Sakai, T. and Nagasawa, T.
Simple, rapid and sensitive determination of plasma taurine by high-performance liquid chromatography using pre-column derivative formation with fluorescamine 576(1992)155
Sallustio, B. C., Morris, R. G. and Horowitz, J. D.
High-performance liquid chromatographic determination of sotalol in plasma. I. Application to the disposition of sotalol enantiomers in humans 576(1992)321
Satoh, N., see Kawakami, Y. 576(1992)63
Schornagel, J. H., see Van Tellingen, O. 576(1992)158
Scott, M. D., Baudendistel, L. J. and Dahms, T. E.
Rapid separation of creatine, phosphocreatine and adenosine metabolites by ion-pair reversed-phase high-performance liquid chromatography in plasma and cardiac tissue 576(1992)149
Seay, S. S., see Tyczkowska, K. L. 576(1992)305
Sertillanges, P., see Rivat, C. 576(1992)71
Shara, M. A., Dickson, P. H., Bagchi, D. and Stohs, S. J.
Excretion of formaldehyde, malondialdehyde, acetaldehyde and acetone in the urine of rats in response to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin, paraquat, endrin and carbon tetrachloride 576(1992)221

Shibasaki, H., Arai, I., Furuta, T. and Kasuya, Y.
Simultaneous determination of cortisol and cortisone in human plasma by stable-isotope dilution mass spectrometry 576(1992)47
Shibasaki, H., see Furuta, T. 576(1992)213
Shimamoto, C., see Kawakami, Y. 576(1992)63
Sips, J. H. M., see Van Tellingen, O. 576(1992)158
Spöndlin, C., see Küsters, E. 576(1992)179
Stein, J., Kulemeier, J., Lembcke, B. and Caspary, W. F.
Simple and rapid method for determination of short-chain fatty acids in biological materials by high-performance liquid chromatography with ultraviolet detection 576(1992)53
Stohs, S. J., see Shara, M. A. 576(1992)221
Stoltz, J.-F., see Regnault, V. 576(1992)87
Stoltz, J.-F., see Rivat, C. 576(1992)71
Storme, G. A., see Van Belle, S. J.-P. 576(1992)351
Stoskopf, M. K., see Tyczkowska, K. L. 576(1992)305
Takagi, T., see Takahara, E. 576(1992)174
Takahara, E., Fukuoka, H., Takagi, T., Nagata, O. and Kato, H.
Simultaneous determination of a new gastrointestinal prokinetic agent (HSR-803) and its metabolites in human serum and urine by high-performance liquid chromatography using automated column-switching 576(1992)174
Tsai, T.-H. and Chen, C.-F.
Determination of glycyrrhizin in rabbit plasma by high-performance liquid chromatography with photodiode-array ultraviolet detection and its pharmacokinetics application 576(1992)170
Tsao, C. S., see Leung, P. Y. 576(1992)245
Tyczkowska, K. L., Seay, S. S., Stoskopf, M. K. and Aucoin, D. P.
Determination of ceftazidime in dolphin serum by liquid chromatography with ultraviolet–visible detection and confirmation by thermospray liquid chromatography–mass spectrometry 576(1992)305
Ubbink, J. B., see Lagendijk, J. 576(1992)95
Vallar, L., see Regnault, V. 576(1992)87
Van Belle, S. J.-P., De Smet, M., Monsaert, C., Geerts, F., Storme, G. A. and Massart, D. L.
High-performance liquid chromatographic determination of navelbine in $MO_4$ mouse fibrosarcoma cells and biological fluids 576(1992)351
Van der Greef, J., see Poon, G. K. 576(1992)235
Van der Hoeven, R. A. M., see Poon, G. K. 576(1992)235
Van Tellingen, O., Sips, J. H. M., Beijnen, J. H., Schornagel, J. H. and Nooyen, W. J.
Bioanalysis of the investigational anti-tumour drug 5,10-dideaza-5,6,7,8-tetrahydrofolic acid by high-performance liquid chromatography with ultraviolet detection 576(1992)158
Vermaak, W. J. H., see Lagendijk, J. 576(1992)95
Visconti, M., see Lombardi, F. 576(1992)129
Wildfeuer, A., see Riedel, K.-D. 576(1992)358
Wyss, R. and Bucheli, F.
Quantitative analysis of retinoids in biological fluids by high-performance liquid chromatography using column switching. III. Determination of the arotinoid sumarotene and its *Z*-isomer in human and animal plasma 576(1992)111
Yoshida, H., see Morita, I. 576(1992)334
Zangirolami, L., see Mariot, R. 576(1992)143
Zimmermann, T., see Riedel, K.-D. 576(1992)358